



Fermi National Accelerator Laboratory

FERMILAB-Conf-95/251-E

D0

Diphoton Production in $p\bar{p}$ Collisions at $\sqrt{s} = 1.8$ TeV

S. Abachi et al.

The D0 Collaboration

*Fermi National Accelerator Laboratory
P.O. Box 500, Batavia, Illinois 60510*

July 1995

Submitted to the *17th International Symposium on Lepton-Photon Interactions (LP95)*,
Beijing, China, August 10-15, 1995

Disclaimer

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, expressed or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Diphoton Production in $p\bar{p}$ Collisions at $\sqrt{s} = 1.8$ TeV

The DØ Collaboration ¹

July 1995

We present measurements of the inclusive $\gamma\gamma$ cross section (as a function of invariant mass and photon E_T), in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV, made using the DØ detector at the Fermilab Tevatron collider. The next to leading order (NLO) QCD prediction is found to be in good agreement with the data. The effects of invariant mass and diphoton balance cuts, which test the next-to-leading order contributions to the cross section, are investigated. We also compare the distribution of k_T between samples of diphotons and highly electromagnetic jets, and find that the NLO QCD prediction models the shape of the $\gamma\gamma$ k_T distribution quite well.

S. Abachi,¹² B. Abbott,³⁴ M. Abolins,²³ B.S. Acharya,⁴¹ I. Adam,¹⁰ D.L. Adams,³⁵ M. Adams,¹⁵ S. Ahn,¹² H. Aihara,²⁰ J. Alitti,³⁷ G. Álvarez,¹⁶ G.A. Alves,⁸ E. Amidi,²⁷ N. Amos,²² E.W. Anderson,¹⁷ S.H. Aronson,³ R. Astur,³⁹ R.E. Avery,²⁹ A. Baden,²¹ V. Balamurali,³⁰ J. Balderston,¹⁴ B. Baldin,¹² J. Bantly,⁴ J.F. Bartlett,¹² K. Bazizi,⁷ J. Bendich,²⁰ S.B. Beri,³² I. Bertram,³⁵ V.A. Bezzubov,³³ P.C. Bhat,¹² V. Bhatnagar,³² M. Bhattacharjee,¹¹ A. Bischoff,⁷ N. Biswas,³⁰ G. Blazey,¹² S. Blessing,¹³ P. Bloom,⁵ A. Boehnlein,¹² N.I. Bojko,³³ F. Borchering,¹² J. Borders,³⁶ C. Boswell,⁷ A. Brandt,¹² R. Brock,²³ A. Bross,¹² D. Buchholz,²⁹ V.S. Burtovoi,³³ J.M. Butler,¹² D. Casey,³⁶ H. Castilla-Valdez,⁹ D. Chakraborty,³⁹ S.-M. Chang,²⁷ S.V. Chekulaev,³³ L.-P. Chen,²⁰ W. Chen,³⁹ L. Chevalier,³⁷ S. Chopra,³² B.C. Choudhary,⁷ J.H. Christenson,¹² M. Chung,¹⁵ D. Claes,³⁹ A.R. Clark,²⁰ W.G. Cobau,²¹ J. Cochran,⁷ W.E. Cooper,¹² C. Cretsinger,³⁶ D. Cullen-Vidal,⁴ M.A.C. Cummings,¹⁴ D. Cutts,⁴ O.I. Dahl,²⁰ K. De,⁴² M. Demarteau,¹² R. Demina,²⁷ K. Denisenko,¹² N. Denisenko,¹² D. Denisov,¹² S.P. Denisov,³³ W. Dharmaratna,¹³ H.T. Diehl,¹² M. Diesburg,¹² G. Di Loreto,²³ R. Dixon,¹² P. Draper,⁴² J. Drinkard,⁶ Y. Ducros,³⁷ S.R. Dugad,⁴¹ S. Durston-Johnson,³⁶ D. Edmunds,²³ J. Ellison,⁷ V.D. Elvira,^{12,†} R. Engemann,³⁹ S. Eno,²¹ G. Eppley,³⁵ P. Ermolov,²⁴ O.V. Eroshin,³³ V.N. Evdokimov,³³ S. Fahey,²³ T. Fahland,⁴ M. Fatyga,³ M.K. Fatyga,³⁶ J. Featherly,³ S. Feher,³⁹ D. Fein,² T. Ferbel,³⁶ G. Finocchiaro,³⁹ H.E. Fisk,¹² Yu. Fisysak,²⁴ E. Flattum,²³ G.E. Forden,² M. Fortner,²⁸ K.C. Frame,²³ P. Franzini,¹⁰ S. Fuess,¹² A.N. Galjaev,³³ E. Gallas,⁴² C.S. Gao,^{12,*} S. Gao,^{12,*} T.L. Geld,²³ R.J. Genik II,²³ K. Genser,¹² C.E. Gerber,^{12,§} B. Gibbard,³ V. Glebov,³⁶ S. Glenn,⁵ B. Gobbi,²⁹ M. Goforth,¹³ A. Goldschmidt,²⁰ B. Gómez,¹ P.I. Goncharov,³³ H. Gordon,³ L.T. Goss,⁴³ N. Graf,³ P.D. Grannis,³⁹ D.R. Green,¹² J. Green,²⁸ H. Greenlee,¹² G. Griffin,⁶ N. Grossman,¹² P. Grudberg,²⁰ S. Grünendahl,³⁶ W. Gu,^{12,*} G. Guglielmo,³¹ J.A. Guida,³⁹ J.M. Guida,³ W. Guryn,³ S.N. Gurzhiev,³³ P. Gutierrez,³¹ Y.E. Gutnikov,³³ N.J. Hadley,²¹ H. Haggerty,¹² S. Hagopian,¹³ V. Hagopian,¹³ K.S. Hahn,³⁶ R.E. Hall,⁶ S. Hansen,¹² R. Hatcher,²³ J.M. Hauptman,¹⁷ D. Hedin,²⁸ A.P. Heinson,⁷ U. Heintz,¹² R. Hernández-Montoya,⁹ T. Heuring,¹³ R. Hirosky,¹³ J.D. Hobbs,¹² B. Hoeneisen,^{1,¶} J.S. Hoftun,⁴ F. Hsieh,²² Ting Hu,³⁹ Tong Hu,¹⁶ T. Huehn,⁷ S. Igarashi,¹² A.S. Ito,¹² E. James,² J. Jaques,³⁰ S.A. Jerger,²³ J.Z.-Y. Jiang,³⁹ T. Joffe-Minor,²⁹ H. Johari,²⁷ K. Johns,² M. Johnson,¹² H. Johnstad,⁴⁰ A. Jonckheere,¹² M. Jones,¹⁴ H. Jöstlein,¹² S.Y. Jun,²⁹ C.K. Jung,³⁹ S. Kahn,³ G. Kalbfleisch,³¹ J.S. Kang,¹⁸ R. Kehoe,³⁰ M.L. Kelly,³⁰ A. Kernan,⁷ L. Kerth,²⁰ C.L. Kim,¹⁸ S.K. Kim,³⁸ A. Klatchko,¹³ B. Klima,¹² B.I. Klockhov,³³

¹Submitted to the XVII International Symposium on Lepton-Photon Interactions (LP95), Beijing, China, August 10–15, 1995.

C. Klopfenstein,³⁹ V.I. Klyukhin,³³ V.I. Kochetkov,³³ J.M. Kohli,³² D. Koltick,³⁴ A.V. Kostritskiy,³³ J. Kotcher,³ J. Kourlas,²⁶ A.V. Kozelov,³³ E.A. Kozlovski,³³ M.R. Krishnaswamy,⁴¹ S. Krzywdzinski,¹² S. Kunori,²¹ S. Lami,³⁹ G. Landsberg,¹² R.E. Lanou,⁴ J-F. Lebrat,³⁷ A. Leflat,²⁴ H. Li,³⁹ J. Li,⁴² Y.K. Li,²⁹ Q.Z. Li-Demarteau,¹² J.G.R. Lima,⁸ D. Lincoln,²² S.L. Linn,¹³ J. Linnemann,²³ R. Lipton,¹² Y.C. Liu,²⁹ F. Lobkowicz,³⁶ S.C. Loken,²⁰ S. Lökös,³⁹ L. Lueking,¹² A.L. Lyon,²¹ A.K.A. Maciel,⁸ R.J. Madaras,²⁰ R. Madden,¹³ I.V. Mandrichenko,³³ Ph. Mangeot,³⁷ S. Mani,⁵ B. Mansoulié,³⁷ H.S. Mao,^{12,*} S. Margulies,¹⁵ R. Markeloff,²⁸ L. Markosky,² T. Marshall,¹⁶ M.I. Martin,¹² M. Marx,³⁹ B. May,²⁹ A.A. Mayorov,³³ R. McCarthy,³⁹ T. McKibben,¹⁵ J. McKinley,²³ T. McMahon,³¹ H.L. Melanson,¹² J.R.T. de Mello Neto,⁸ K.W. Merritt,¹² H. Miettinen,³⁵ A. Milder,² A. Mincer,²⁶ J.M. de Miranda,⁸ C.S. Mishra,¹² M. Mohammadi-Baarmand,³⁹ N. Mokhov,¹² N.K. Mondal,⁴¹ H.E. Montgomery,¹² P. Mooney,¹ M. Mudan,²⁶ C. Murphy,¹⁶ C.T. Murphy,¹² F. Nang,⁴ M. Narain,¹² V.S. Narasimham,⁴¹ A. Narayanan,² H.A. Neal,²² J.P. Negret,¹ E. Neis,²² P. Nemethy,²⁶ D. Nešić,⁴ D. Norman,⁴³ L. Oesch,²² V. Oguri,⁸ E. Oltman,²⁰ N. Oshima,¹² D. Owen,²³ P. Padley,³⁵ M. Pang,¹⁷ A. Para,¹² C.H. Park,¹² Y.M. Park,¹⁹ R. Partridge,⁴ N. Parua,⁴¹ M. Paterno,³⁶ J. Perkins,⁴² A. Peryshkin,¹² M. Peters,¹⁴ H. Piekarz,¹³ Y. Pischalnikov,³⁴ A. Pluquet,³⁷ V.M. Podstavkov,³³ B.G. Pope,²³ H.B. Prosper,¹³ S. Protopopescu,³ D. Pušeljčić,²⁰ J. Qian,²² P.Z. Quintas,¹² R. Raja,¹² S. Rajagopalan,³⁹ O. Ramirez,¹⁵ M.V.S. Rao,⁴¹ P.A. Rapidis,¹² L. Rasmussen,³⁹ A.L. Read,¹² S. Reucroft,²⁷ M. Rijssenbeek,³⁹ T. Rockwell,²³ N.A. Roe,²⁰ P. Rubinov,³⁹ R. Ruchti,³⁰ S. Rusin,²⁴ J. Rutherford,² A. Santoro,⁸ L. Sawyer,⁴² R.D. Schamberger,³⁹ H. Schellman,²⁹ J. Sculli,²⁶ E. Shabalina,²⁴ C. Shaffer,¹³ H.C. Shankar,⁴¹ R.K. Shivpuri,¹¹ M. Shupe,² J.B. Singh,³² V. Sirotenko,²⁸ W. Smart,¹² A. Smith,² R.P. Smith,¹² R. Snihur,²⁹ G.R. Snow,²⁵ S. Snyder,³⁹ J. Solomon,¹⁵ P.M. Sood,³² M. Sosebee,⁴² M. Souza,⁸ A.L. Spadafora,²⁰ R.W. Stephens,⁴² M.L. Stevenson,²⁰ D. Stewart,²² D.A. Stoianova,³³ D. Stoker,⁶ K. Streets,²⁶ M. Strovink,²⁰ A. Taketani,¹² P. Tamburello,²¹ J. Tarazi,⁶ M. Tartaglia,¹² T.L. Taylor,²⁹ J. Teiger,³⁷ J. Thompson,²¹ T.G. Trippe,²⁰ P.M. Tuts,¹⁰ N. Varelas,²³ E.W. Varnes,²⁰ P.R.G. Virador,²⁰ D. Vititoe,² A.A. Volkov,³³ A.P. Vorobiev,³³ H.D. Wahl,¹³ G. Wang,¹³ J. Wang,^{12,*} L.Z. Wang,^{12,*} J. Warchol,³⁰ M. Wayne,³⁰ H. Weerts,²³ F. Wen,¹³ W.A. Wenzel,²⁰ A. White,⁴² J.T. White,⁴³ J.A. Wightman,¹⁷ J. Wilcox,²⁷ S. Willis,²⁸ S.J. Wimpenny,⁷ J.V.D. Wirjawan,⁴³ J. Womersley,¹² E. Won,³⁶ D.R. Wood,¹² H. Xu,⁴ R. Yamada,¹² P. Yamin,³ C. Yanagisawa,³⁹ J. Yang,²⁶ T. Yasuda,²⁷ C. Yoshikawa,¹⁴ S. Youssef,¹³ J. Yu,³⁶ Y. Yu,³⁸ Y. Zhang,^{12,*} Y.H. Zhou,^{12,*} Q. Zhu,²⁶ Y.S. Zhu,^{12,*} Z.H. Zhu,³⁶ D. Zieminska,¹⁶ A. Zieminski,¹⁶ and A. Zylberstein³⁷

¹Universidad de los Andes, Bogotá, Colombia

²University of Arizona, Tucson, Arizona 85721

³Brookhaven National Laboratory, Upton, New York 11973

⁴Brown University, Providence, Rhode Island 02912

⁵University of California, Davis, California 95616

⁶University of California, Irvine, California 92717

⁷University of California, Riverside, California 92521

⁸LAFEX, Centro Brasileiro de Pesquisas Físicas, Rio de Janeiro, Brazil

⁹CINVESTAV, Mexico City, Mexico

¹⁰Columbia University, New York, New York 10027

¹¹Delhi University, Delhi, India 110007

¹²Fermi National Accelerator Laboratory, Batavia, Illinois 60510

¹³Florida State University, Tallahassee, Florida 32306

¹⁴University of Hawaii, Honolulu, Hawaii 96822

¹⁵University of Illinois at Chicago, Chicago, Illinois 60607

¹⁶Indiana University, Bloomington, Indiana 47405

¹⁷Iowa State University, Ames, Iowa 50011

¹⁸Korea University, Seoul, Korea

¹⁹Kyungshung University, Pusan, Korea

²⁰Lawrence Berkeley Laboratory and University of California, Berkeley, California 94720

²¹University of Maryland, College Park, Maryland 20742

²²University of Michigan, Ann Arbor, Michigan 48109

²³Michigan State University, East Lansing, Michigan 48824

²⁴Moscow State University, Moscow, Russia

²⁵University of Nebraska, Lincoln, Nebraska 68588

²⁶New York University, New York, New York 10003

²⁷Northeastern University, Boston, Massachusetts 02115

- ²⁸Northern Illinois University, DeKalb, Illinois 60115
²⁹Northwestern University, Evanston, Illinois 60208
³⁰University of Notre Dame, Notre Dame, Indiana 46556
³¹University of Oklahoma, Norman, Oklahoma 73019
³²University of Panjab, Chandigarh 16-00-14, India
³³Institute for High Energy Physics, 142-284 Protvino, Russia
³⁴Purdue University, West Lafayette, Indiana 47907
³⁵Rice University, Houston, Texas 77251
³⁶University of Rochester, Rochester, New York 14627
³⁷CEA, DAPNIA/Service de Physique des Particules, CE-SACLAY, France
³⁸Seoul National University, Seoul, Korea
³⁹State University of New York, Stony Brook, New York 11794
⁴⁰SSC Laboratory, Dallas, Texas 75237
⁴¹Tata Institute of Fundamental Research, Colaba, Bombay 400005, India
⁴²University of Texas, Arlington, Texas 76019
⁴³Texas A&M University, College Station, Texas 77843

INTRODUCTION

Diphoton production at the Tevatron is of interest, firstly, as a test of QCD. Next to leading order predictions are available, and CDF have reported [1] a cross section about three times higher than expected (though more recent unpublished results [2] are lower). Secondly, it is a major irreducible background to Higgs discovery in the channel $H \rightarrow \gamma\gamma$ at the LHC. Thirdly it provides a way to test recent suggestions [3] that significant additional k_T (due to soft gluon radiation) needs to be added to perturbative QCD calculations in order to correctly model photon production.

This study uses 60 pb^{-1} of data taken during 1994–95 with the DØ detector. Events were recorded if they satisfied the diphoton trigger, which required two electromagnetic (EM) calorimeter towers above threshold at the hardware trigger level, and two EM clusters with $E_T > 12 \text{ GeV}$ at the software trigger level.

DATA SELECTION

Events were required to have two photon candidates found in the DØ calorimeter which satisfied the following cuts:

- $E_T^1 > 20 \text{ GeV}$, $E_T^2 > 18 \text{ GeV}$;
- $|\eta| < 0.9$;
- Isolation with $E_T^{R=0.4} - E_T^{R=0.2} < 2 \text{ GeV}$, where $E_T^{R=0.4}(E_T^{R=0.2})$ is the energy contained in a cone of radius $R = \sqrt{\Delta\eta^2 + \Delta\phi^2} = 0.4(0.2)$ about the direction of the photon;
- Fraction of the cluster energy in the EM calorimeter (EM fraction) > 0.96 ;
- Have a shower shape consistent with a test beam electron;
- Not be close to an azimuthal calorimeter crack (10% of 2π was excluded);
- No track is found in the road in front of the EM cluster.

The combined acceptance and efficiency of these cuts is estimated to be 0.28 ± 0.03 from detailed Monte Carlo simulations including noise and pileup effects [4]. This includes a detector acceptance 0.81 ± 0.01 , a combined

trigger and cut efficiency of 0.64 ± 0.03 , and a probability that no track from the underlying event be found in the road of 0.925 ± 0.05 . (The latter is estimated from data by counting the number of tracks found in random roads in $Z \rightarrow ee$ events). The efficiencies were checked against data using $Z \rightarrow ee$ events in the region $20 < p_T^e < 50$ GeV/c and found to agree to within 4% [4]; this 4% is included in the error on the efficiency estimation.

In addition to the cuts outlined above, it was found necessary to impose an invariant mass requirement to remove $Z \rightarrow ee$ events where both electron tracks were lost because of tracking inefficiencies. (The number of observed events in the Z mass region is consistent with a tracking efficiency of about 0.9). Events with $80 < m_{\gamma\gamma} < 110$ GeV were therefore excluded.

217 events remain after these selections.

A complementary sample of highly electromagnetic jets was selected from the same diphoton trigger and dataset. The EM jets were selected by requiring the same kinematic cuts as above, but:

- Anti-isolation with $E_T^{R=0.4} - E_T^{R=0.2} \geq 2$ GeV;
- More than 1% of the shower energy in the first layer ($2X_0$) of the EM calorimeter;
- EM fraction > 0.90 ;
- One or more of: EM fraction < 0.96 , shower shape not consistent with electron, or ≥ 2 tracks found in the road in front of the EM cluster.

These selections yield 81 events. This sample is expected to be composed almost entirely of dijet events where both jets have fragmented into electromagnetically-decaying particles. No real isolated photons are expected to remain in this sample.

BACKGROUND ESTIMATION

Even after the photon selection cuts described above, a significant background still remains from events where a jet has fragmented into a hard π^0 or η meson and which is found in the detector as a photon. The probability for this to occur (given the DØ photon requirements) is $\sim 5 \times 10^{-4}$, but since the QCD dijet cross section is a few $\times 10^6$ times higher than the $\gamma\gamma$ cross section, and the $\gamma + jet$ cross section is a few $\times 10^3$ times higher than the $\gamma\gamma$ cross section, both of these must be considered as potential sources of background.

The background estimation technique relies on the fraction of photon candidates having an energy in the first layer of the EM calorimeter ($2X_0$) which is less than 0.01 of the shower energy ($EM1/E < 0.01$) as a discriminant. Real photons have a higher probability to have $EM1/E < 0.01$ than the multiphoton backgrounds from π^0 and η decays, because they are less likely to convert early. (The $EM1/E$ distribution has been shown to be stable as a function of luminosity.) The probabilities ϵ_γ and ϵ_{jet} for photons and electromagnetically-fragmenting jets, respectively, to have $EM1/E < 0.01$, are estimated from detailed Monte Carlo [4] (jets are treated as an admixture of π^0 and η mesons). Both ϵ_γ and ϵ_{jet} depend upon the photon E_T , but given the very limited statistics, it is not possible to estimate the background as a function of E_T ; instead ϵ_γ and ϵ_{jet} have been evaluated at $E_T = 29$ GeV, which is the mean E_T of the photons in the signal sample and of the jets in the background sample. Here $\epsilon_\gamma = 0.231$ and $\epsilon_{jet} = 0.084$.

We may then write:

$$\begin{pmatrix} N_{PP} \\ N_{PF} \\ N_{FF} \end{pmatrix} = \begin{pmatrix} \epsilon_\gamma^2 & \epsilon_\gamma \epsilon_{jet} & \epsilon_{jet}^2 \\ 2\epsilon_\gamma(1-\epsilon_\gamma) & \epsilon_\gamma(1-\epsilon_{jet}) + \epsilon_{jet}(1-\epsilon_\gamma) & 2\epsilon_{jet}(1-\epsilon_{jet}) \\ (1-\epsilon_\gamma)^2 & (1-\epsilon_\gamma)(1-\epsilon_{jet}) & (1-\epsilon_{jet})^2 \end{pmatrix} \begin{pmatrix} N_{\gamma\gamma} \\ N_{\gamma j} \\ N_{jj} \end{pmatrix} \quad (1)$$

where (N_{PP}, N_{PF}, N_{FF}) are the numbers of events with (2,1,0) photons satisfying $EM1/E < 0.01$, and $(N_{\gamma\gamma}, N_{\gamma j}, N_{jj})$ are the numbers of events which are true diphotons, photon+jet and dijets. We may then obtain

18/07/95 17.00




FIG. 1. Diphoton differential cross section $d\sigma/dm_{\gamma\gamma}$ as a function of the diphoton invariant mass $m_{\gamma\gamma}$. The plot contains 217 candidate events with an expected purity of 0.27.

$(N_{\gamma\gamma}, N_{\gamma j}, N_{jj})$ by matrix inversion. For $(N_{PP}, N_{PF}, N_{FF}) = (7, 49, 161)$ this inversion predicts an unphysical solution (the number of $\gamma + jet$ events in the sample is negative). The unphysical solution probably reflects the rather large statistical uncertainties on the observed numbers of events, particularly N_{PP} .

We therefore introduce an additional constraint on the relative amount of the two backgrounds, $N_{\gamma j}/(N_{\gamma j} + N_{jj})$. The $\gamma + jet$ background may be estimated as $N_{\gamma j} = \mathcal{L}\sigma_{\gamma j} \times P$ where P is the probability for a jet to fluctuate into a photon candidate. Similarly, $N_{jj} = \mathcal{L}\sigma_{u\gamma jj} \times P$, where $\sigma_{u\gamma jj}$ is the cross section for fake photons. Then $N_{\gamma j}/(N_{\gamma j} + N_{jj}) = \sigma_{\gamma j}/(\sigma_{\gamma j} + \sigma_{u\gamma jj})$. From the single inclusive photon analysis, this ratio is known; it is the purity of the photon candidate sample. For $E_T = 29$ GeV, we find [4] $\sigma_{\gamma j}/(\sigma_{\gamma j} + \sigma_{u\gamma jj}) = 0.31 \pm 0.05$. With $N_{\gamma j}/(N_{\gamma j} + N_{jj})$ fixed to this value, the 3×3 matrix equation can be reduced to 2 equations in 2 unknowns and solved directly. The variable N_{PP} is not used in the solution (as it has the largest uncertainty). We obtain $f = 0.268 \pm 0.15 \pm 0.025$, where the first error is statistical and the second reflects the uncertainty in $N_{\gamma j}/(N_{\gamma j} + N_{jj})$. The inverted equations predict $N_{PP} = 4$, which is in reasonable agreement with the observed value.

In order to estimate the systematic error that may arise from estimating the background using ϵ_γ and ϵ_{jet} values which are constant rather than varying with E_T , ϵ_γ and ϵ_{jet} were varied in a correlated way by amounts corresponding to $\pm 1\sigma$ in the E_T distribution. This results in an additional error of $(+0.17, -0.14)$ on the $\gamma\gamma$ signal purity.

The systematic errors on the cross section have been combined in quadrature. They include $\pm 9.6\%$ from acceptance and efficiency, $\pm 5\%$ luminosity uncertainty, $\pm 56\%$ statistical error on the background subtraction, $(+63\%, -52\%)$ systematic error on the background subtraction from varying E_T and $\pm 9.3\%$ from varying $N_{\gamma j}/(N_{\gamma j} + N_{jj})$. The combined error is $(+85\%, -75\%)$.




FIG. 2. Diphoton differential cross section $d\sigma\epsilon/dE_T^\gamma$ as a function of the photon transverse momentum E_T^γ . The probability to pass the invariant mass cut, ϵ , has not been corrected for. The plot contains 217 candidate events with an expected purity of 0.27.

The background subtraction has been carried out by multiplying the $\gamma\gamma$ candidate distribution by the signal purity. This is justified by the observation that the $\gamma\gamma$ candidates and the EM jet sample have very similar invariant mass and p_T distributions, so the background contribution to all bins is expected to be similar.

DIPHOTON CROSS SECTION

Fig. 1 shows the differential diphoton cross section $d\sigma/dm_{\gamma\gamma}$ as a function of the diphoton invariant mass $m_{\gamma\gamma}$. The data are compared with a next-to-leading order (NLO) QCD prediction [5], evaluated at a renormalization scale $\mu = E_T$ with CTEQ3M parton distributions; the data and theory are in good agreement over the whole range of invariant mass.

Fig. 2 shows the differential diphoton cross section $d\sigma\epsilon/dE_T^\gamma$ as a function of the photon transverse energy E_T^γ (two entries per event). The dip in the cross section around $E_T^\gamma = 50$ GeV is due to the invariant mass cut applied to remove Z events; we have not corrected for the probability to pass this cut (ϵ). The data are again compared with the next-to-leading order QCD prediction [5], and the agreement between data and theory is once again good, even in the region affected by the invariant mass cut. It is interesting to note that at leading order, there would be no events at all in this region, since the whole leading order cross section is due to back-to-back photon pairs. The population of events here is then a test of the next to leading order contributions to the QCD cross section. We may explore this effect further by introducing a cut on the imbalance between the two photons, z :

18/07/95 17.00

18/07/95 17.00




FIG. 3. Diphoton differential cross section $d\sigma/dE_T^\gamma$ as a function of the photon transverse momentum E_T^γ , for imbalance cuts of (a) $z \geq 0.5$ and (b) $z \geq 0.8$. The probability to pass the invariant mass cut, ϵ , has not been corrected for.

$$z = \frac{|\mathbf{p}_1 \cdot \mathbf{p}_2|}{\mathbf{p}_1^2} \quad (2)$$

where \mathbf{p}_1 and \mathbf{p}_2 are the vector momenta of the two photons. Requiring $z \geq 0.5$ or $z \geq 0.8$ selects events with increasingly back-to-back topologies, and correspondingly restricts the cross section to its leading order part, as can be seen in Fig. 3. The NLO QCD prediction remains in good agreement with the data for both of these cases.

DIPHOTON K_T

We define the diphoton k_T by:

$$k_T = |\mathbf{p}_T^1 + \mathbf{p}_T^2| \quad (3)$$

where \mathbf{p}_T^1 and \mathbf{p}_T^2 are the vector transverse momenta of the two photons. Because $k_T \ll p_T^{1,2}$, it is not amenable to perturbative calculation, and *ad hoc* models of soft gluon radiation have been proposed to predict the distribution.

If soft gluon radiation is indeed important, then we may expect that the k_T distribution will be broader for dijet events (predominantly $gg \rightarrow gg$ scattering at the Tevatron) than for diphotons. This is because the probability for radiation is higher off gluon lines, and final state radiation can also contribute in this case.

Because we wish to investigate differences between the k_T distributions for diphoton and dijet events, it is desirable to use the purest possible $\gamma\gamma$ sample, and so an additional requirement was made: that one or both photons have $EM1/E < 0.01$. This results in 57 events with an estimated purity of 0.41. This enriched $\gamma\gamma$ sample will be compared with the EM jet sample. Since the purity of the signal sample is known, the background sample may be directly subtracted to yield a background-subtracted $\gamma\gamma$ distribution. (This assumes, of course, that the k_T of the $\gamma + jet$ and dijet contributions to the background are similar).

The normalized k_T distributions $1/N dN/dk_T$ for the three cases are shown in Fig. 4. While the general shape of both the diphoton and dijet distributions is similar, the dijet sample has a higher mean and most probable k_T (the fraction of events with $k_T < 4 \text{ GeV}$ is $25 \pm 6\%$ for the diphotons and $14 \pm 4\%$ for the dijets). However,




FIG. 4. Normalized differential k_T distributions, $1/N dN/dk_T$, for (a) an enriched sample of $\gamma\gamma$ candidates (56 events with a purity of 0.41), (b) highly electromagnetic jets (81 events), and (c) the $\gamma\gamma$ sample after subtraction of the background distribution.

it must be noted that the EM jet selection permits more hadronic energy in the clusters than for the photon cuts, so the dijet events will have a worse energy resolution. A simple estimate suggests that for $p_T = 30$ GeV candidates at $\eta = 0$, the k_T resolution is ~ 1.2 GeV for diphotons and ~ 1.7 GeV for EM jets. This could account for some or all of the difference in k_T distributions; more detailed modelling will be required to fully understand the magnitude of this effect. We cannot therefore claim at this time that the observed difference in k_T between diphotons and dijets has its origin in physics rather than detector effects.

The NLO QCD prediction [5] is shown superimposed on the the background-subtracted $\gamma\gamma$ distribution. The theory Monte Carlo included smearing the photon 4-vectors according to the $0.15\sqrt{E}$ energy resolution of the DØ calorimeter. The NLO QCD prediction models the k_T distribution quite well, even though it has been claimed [3] that perturbative QCD should not be able to match the low- k_T behavior (below about 5 GeV) correctly.

CONCLUSIONS

We have measured the inclusive cross section $p\bar{p} \rightarrow \gamma\gamma$ as a function of invariant mass and photon E_T . The next to leading order QCD prediction is in good agreement with the data. It is also able to correctly model the effect of invariant mass and momentum imbalance (z) cuts imposed on the data, which test the next-to-leading order contributions to the cross section.

The distribution of k_T has been compared between a sample of diphoton candidates and highly electromagnetic jets. The two distributions are generally similar, but the jet sample has a higher mean and most probable k_T . This may merely indicate the effects of worse energy resolution. A background-subtracted diphoton sample is compared with the NLO QCD prediction for k_T . The theoretical prediction matches the observed distribution quite well, although it is not expected to model the data below $k_T \sim 5\text{GeV}$.

ACKNOWLEDGMENTS

We thank the Fermilab Accelerator, Computing, and Research Divisions, and the support staffs at the collaborating institutions for their contributions to the success of this work. We also acknowledge the support of the U.S. Department of Energy, the U.S. National Science Foundation, the Commissariat à l'Energie Atomique in France, the Ministry for Atomic Energy and the Ministry of Science and Technology Policy in Russia, CNPq in Brazil, the Departments of Atomic Energy and Science and Education in India, Colciencias in Colombia, CONACyT in Mexico, the Ministry of Education, Research Foundation and KOSEF in Korea, and the A.P. Sloan Foundation.

REFERENCES

- * Visitor from IHEP, Beijing, China.
 - † Visitor from CONICET, Argentina.
 - § Visitor from Universidad de Buenos Aires, Argentina.
 - ¶ Visitor from Univ. San Francisco de Quito, Ecuador.
1. CDF Collaboration, Phys. Rev. Lett. **70**, 2232.
 2. R. Blair, at the 10th Topical Workshop on $p\bar{p}$ Collider Physics, Fermilab, May 1995.
 3. J. Huston *et al.*, 'A Global Study of Direct Photon Production,' MSU-HEP-41027, CTEQ-407, January 1995.
 4. S. Fahey, Ph.D. Thesis, Michigan State University, July 1995.
 5. B. Bailey, private communications.